


Clinical & Counselling Psychology Review (CCPR)

Volume 7 Issue 2, Fall 2025

ISSN_(P): 2412-5253, ISSN_(E): 2706-8676

Homepage: <https://journals.umt.edu.pk/index.php/CCPR>



Title: Efficacy of Positive Psychotherapy (PPT) in Reducing the Symptoms of Depression and Improving Life Satisfaction among University Students

Author (s): Amina Shahzadi and Faiza Safdar

Affiliation (s): University of the Punjab, Lahore, Pakistan

DOI: <https://doi.org/10.32350/ccpr.72.03>

History Received: August 26, 2025, Revised: November 06, 2025, Accepted: November 18, 2025, Published: December 31, 2025

Citation: Shahzadi, A., & Safdar, F. (2025). Efficacy of positive psychotherapy (PPT) in reducing the symptoms of depression and improving life satisfaction among university students. *Clinical and Counselling Psychology Review*, 7(2), 47–70. <https://doi.org/10.32350/ccpr.72.03>

Copyright: © The Authors

Licensing:  This article is open access and is distributed under the terms of [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

Conflict of Interest: Author(s) declared no conflict of interest



A publication of
Department of Clinical Psychology
University of Management and Technology, Lahore, Pakistan

Efficacy of Positive Psychotherapy (PPT) in Reducing the Symptoms of Depression and Improving Life Satisfaction among University Students

Amina Shahzadi* and Faiza Safdar

Centre for Clinical Psychology, University of the Punjab, Lahore, Pakistan

Abstract

Positive psychotherapy (PPT) is considered one of the optimal therapeutic strategies in the management of depression for contributing to positive emotions, engagement in life, improving interpersonal relationships, and adding meaning in life. The present study aimed to evaluate the efficacy of PPT in reducing the symptoms of depression among university students. In addition, it also examined the role of PPT in enhancing life satisfaction among students. For this, a sample of 22 undergraduate students was taken including students who appeared in the entrance exam (MDCAT) for admission in medical colleges but failed to get in. All these students were assessed using Patient Health Questionnaire (PHQ-9) and Multidimensional Students' Life Satisfaction Scale and met the criteria of depression during their initial assessment. They were divided in two groups i.e., control (no-treatment) and experimental group. The experimental group received six group-sessions of PPT, planned considering Seligman et al. (2006)'s work. At post-intervention level, the assessment was replicated for both groups in order to examine the group differences. The findings revealed that PPT remained effective in reducing depressive symptomatology and improving life satisfaction among university students in five major domains including satisfaction with self, family, educational institution, friends, and living environment. The study was concluded to be a significant contribution in the literature, encouraging the use of PPT in clinical settings. The study findings highlight the efficacy of brief third wave therapeutic interventions in promoting mental health and well-being over traditional therapies.

Keywords: depression, life satisfaction, positive psychotherapy, students

Introduction

Mental health has always been a public health concern which is particularly

*Corresponding Author: amina.cpsy@gmail.com

challenging in low- and middle-income countries (LMICs) including Pakistan (Albarqi et al., 2024). Evidence in this reference, retrieved from Global Burden of Disease Study data for 1990–2019, showed a significantly increased mental health burden in Pakistan over three decades. It has also been observed that depressive disorders are most commonly occurring and contribute significantly to disability-adjusted life-years, demanding the attention of mental health stakeholders (Alvi et al., 2023).

With a growing concern surrounding mental health, the young population has been a focus of research over the past few years across the globe. These years are crucial for education, employment, and social relationships (National Academies of Sciences, Engineering, and Medicine [NASEM], 2019). Evidence suggests that depression and anxiety are among the common mental health conditions experienced by adolescents globally (Silva et al., 2020). It has been estimated that 24 to 34% university students experience depressive symptoms across the globe (Akhtar et al., 2020; Tam et al., 2019). Despite the current global interest in the mental health of the young population, studies from LMICs, with Pakistan being no exception, are scarce (Khan et al., 2021).

It has been documented that approximately 29% of the Pakistani population consists of young adults (15–29 years) (Khan et al., 2021), which is also the age where the onset of depression is reported to be the highest (Substance Abuse and Mental Health Services Administration [SAMHSA], 2022). According to a systematic review, about 43% of university students in Pakistan have been experiencing symptoms of depression, which is alarming (Khan et al., 2021). Likewise, a survey reported the prevalence of depression among 53% university students, followed by the prevalence of anxiety in 43% (Muneer et al., 2024). Recent cross-sectional data obtained from the general population have suggested that almost 20% of the young population in Pakistan suffers from mood disorders, with depressive disorders being highly reported (Rahman et al., 2024). A similar study with young adults has also reported that about 40% of this population in Pakistan is suffering from depression (Ullah et al., 2022).

For this population, the chance to experience psychopathology is particularly higher in the presence of risk factors, which could be biological, psychological, social, academic, or related to lifestyle (Mofatteh, 2020). Evidence suggests that the high prevalence of depression indicates it as a common risk factor for reduced satisfaction with life (Lukaschek et al.,

[2017](#)), with the young student population being no exception (Bukhari & Saba, [2017](#)). Satisfaction with life is also considered an important predictor of depression, anxiety (Liu & Wang, [2024](#)), and positive affect (Saldivia et al., [2023](#)). Data obtained from studies based on students suggest that depression and anxiety negatively predict students' satisfaction with life (Liu & Wang, [2024](#)), while students with a higher level of satisfaction with life score low on depression and anxiety (Ooi et al., [2022](#)). In addition to reduced satisfaction, these mental health conditions can predispose to severe consequences. For example, suicidal behaviors among students (Kabir et al., [2024](#)) particularly after experiencing distress (Bilal & Riaz, [2020](#)). Depression, in particular, is a potential threat to the physical as well as the mental well-being of the students (Irie & Yokomitsu, [2019](#)). Taken together, evidence suggests that depression and other common mental health conditions, such as anxiety and stress, among students often impact their mental well-being and academic performance (Storrie et al., [2010](#)), which warrants the attention of academic and mental health stakeholders.

Among many academic stressors experienced by students, failure to achieve an academic goal can be a major source of distress. A single experience of failure can result in extreme emotional disturbance (Johnson et al., [2017](#)). Considering the Pakistani context, the mental health of students who failed to secure admission to a medical college due to lower scores on the entrance exam for admission in medical colleges is a burning issue (Bilal & Riaz, [2020](#); Fatima et al., [2021](#)). The field of medicine is highly regarded in Pakistan, however, the positions in medical colleges are limited, which makes it challenging for aspirants to get through the exam and secure admission (Bilal & Riaz, [2020](#)). Therefore, one of the potential reasons for poor exam performance or failure could be the pressure or distress experienced by the students, as in the case of students taking MDCAT (Bilal & Riaz, [2020](#)). This distress can significantly affect their academic performance and manifest itself in the form of mental health issues like depression (Deng et al., [2022](#)), which is a commonly experienced issue among students. Evidence suggests that students often experience depression due to emotional disturbance resulting from poor academic outcomes, failures, worries about the future, and career related concerns (Erschens et al., [2018](#); Feng et al., [2022](#); Mathew et al., [2015](#); Moir et al., [2018](#); Sinval et al., [2024](#); Yousef et al., [2017](#)). Factors like failure to achieve a goal (e.g., to get admission in MBBS/BDS due to poor MDCAT performance), subject selection choice, or concerns related to career,

therefore, can be extremely distressing for students and may lead to depression, which is supported by available literature (Erschens et al., 2018; Fatima et al., 2021; Yousef et al., 2017).

In response to the higher prevalence of depression among the student population, increasing efforts to detect and treat depression among students have been carried out with the aim of societal benefits. These efforts aim to control the outcomes of depression, for instance, higher academic withdrawals (Cuijpers et al., 2016). For the treatment or management of depression, one of the important questions is the availability of evidence-based psychotherapies. For this, several evidence-based therapeutic modalities are available; many of these therapies are typically based on cognitive approaches, while some incorporate multiple orientations. Although the intent is generally to implement a therapy across multiple settings, it is important to consider the therapeutic orientation in terms of its applicability to the patient as well as the provider (Cook et al., 2017). In this context, a lack of supporting evidence exists regarding the utility of evidence-based psychotherapies with the youth or student populations, as they may respond differently to the routine treatment (Barnett et al., 2022; Cuijpers et al., 2016). This could be particularly challenging in LMICs where accessibility to the mental health services is a challenge in itself. These challenges may be related to various factors, including financial resources, organization and planning, policies, access and availability of appropriately adapted evidence-based interventions and training. Such factors are influenced by a specific population's cultural belief system and worldview regarding mental health and illness. This interplay may serve as both a strength and a weakness, yet it undeniably affects service utilization (Rathod et al., 2017), in conjunction with mental health stigma (Ahmad & Koncsol, 2022).

Considering the need to address this issue, context, and functions of psychological phenomena of depression, the present study provides evidence on the utility of positive psychotherapy (PPT) for the student population suffering from depression. Theoretically based on the Seligman's positive psychology, positive psychotherapy (PPT) emphasizes that constructing and strengthening positive emotions, meaning (i.e., belonging to and serving something greater than the self), and engagement in life can undo psychopathology (Seligman, 2002). It assumes that an individual has an inherent tendency to seek growth, fulfillment, and

happiness. An effective therapeutic relationship can be developed through the manifestation of these positive resources that are as real as the symptoms of psychopathology (Seligman, [2011](#)). By focusing on both strengths and weaknesses of an individual, positive psychotherapy (PPT) offers an incremental change to the deficit-oriented approach of traditional therapies. The fundamental aim of this therapeutic approach is to increase positive affect, meaning in life, and personal engagement (Rashid, [2014](#)). In this regard, PPT appears to be a useful modality for the Pakistani collectivistic cultural context which emphasizes values, connection, engagement, and meaningfulness in relationships (Ford et al., [2015](#); Liu et al., [2021](#); Memon et al., [2021](#)).

Literature provides significant evidence supporting the effectiveness of positive psychotherapy (PPT) in reducing the symptoms of depression, and increasing the life satisfaction (Asl et al., [2016](#)) with a comparable efficacy to CBT (Hoppen & Morina, [2021](#); Seligman et al., [2006](#)). In fact, evidence is available that shows PPT is more effective in treating depression in comparison to CBT (Asgharipoor et al., [2012](#)). Additionally, delving on the theoretical evidence, PPT seems to be promising in improving life satisfaction. As suggested by Fredrickson's ([2004](#)) broaden-and-build theory, positive emotions broaden an individual's attention and thinking style resulting in flexibility and creativity. PPT enhances positive emotions that broaden cognition and build psychological resources. As a result, individuals are more likely to appraise their lives positively and report greater life satisfaction.

Although the efficacy of PPT has been mainly investigated in terms of treating depression, there are some studies that target other clinical populations. For instance, it has been evaluated for individuals with psychosis (Schrank et al., [2015](#)), anxiety (Goodwin, [2010](#)), borderline personality (Uliaszek et al., [2016](#)), breast cancer (Dowlatabadi et al., [2016](#)), and habitual smoking (Kahler et al., [2014](#)). In the indigenous Pakistani context, there exists a lack of studies on PPT; only one published study could be found that examines the efficacy of PPT with adult females who had mild to moderate depression. Their study reported that PPT was effective in reducing depression and improving happiness (Mazhar & Riaz, [2020](#)). In conclusion, these findings support the assumption that PPT leads to an enhancement of happiness, positive emotions, and well-being, and concurrently results in a decrease in psychopathology. The application of

PPT in interventional research is heterogeneous in terms of both the modifications to the original manual as well as the conditions targeted by PPT as intended by the PPT developers (Walsh et al., [2017](#)).


Method

Study Design

The present study was carried out between Jan-July, 2021, and used a quasi-experimental design to test the efficacy of PPT. The study participants were divided into treatment and control group without any randomization.

Figure 1

CONSORT Flow Diagram



Study Participants

The study participants were approached online through advertisement on social media platforms, and the researchers targeted four high merit departments of the University of the Punjab. Using purposive sampling strategy, students were initially screened for inclusion criteria using a Google Form. Among the 51 initially approached students, 22 participants were enrolled in the study and were evenly classified in the treatment ($n = 11$) and control ($n = 11$) groups (see Figure 1).

The participants who studied pre-medical at college, recently appeared in the entrance exam of medical colleges (MDCAT) for admission in MBBS/BDS but couldn't get admission due to their lower aggregate, and had mild to moderate levels of depression were included. All these participants were ≥ 18 years and currently enrolled as regular students in their first year of university. Those with age above 23 years, who had appeared in the entrance exam twice or more, were undergoing treatment for depression, or had disability or co-morbid psychological disturbance were excluded.

Assessment Measures

Participant Information Sheet

A participant sheet was used to gather personal and demographic information such as age, gender, education, socioeconomic status, family, medical college entrance exam, physical or psychological issues, etc.

Patient Health Questionnaire

The PHQ-9 was used for a quick assessment of depression and its severity. It consisted of 9 items, which were rated on a "0" (*not at all*) to "3" (*nearly every day*) Likert scale. The score on PHQ-9 ranged from 0 to 27, with a diagnostic cutoff being 10. The score on this questionnaire, computed by adding individual item scores, provided information regarding the different severity levels of depression that ranged from minimal to severe. Evidence suggests that PHQ-9 is a psychometrically sound tool for the assessment of depression across populations (Kim & Lee, [2019](#); Kroenke et al., [2001](#)). In the present study, it has shown an alpha value of .86 proving it to be a reliable measure.

Multidimensional Students' Life Satisfaction Scale

The MSLLS was used to assess the participants' level of satisfaction with life. This scale consists of 40 items divided into five subscales tapping satisfaction with life: Self, school, friends, family, and living environment. Each item is rated on a Likert scale marked from "1" (*strongly disagree*) to "6" (*strongly agree*). The scoring of each subscale is done by summing individual item ratings in a subscale and then dividing the sum by the number of items in that subscale. The MSLLS has been evident as a valid and reliable measure of assessment (Huebner, [2001](#); Schnettler et al., [2017](#)). In the present study, the scale proved to be a reliable measure with an alpha value of .90. The alpha values for individual subscales were observed as the following: Family (.87); Friends (.84), School (.89), Environment (.84), and Self (.85).

Treatment Protocol

The therapeutic protocol used was PPT (Rashid & Seligman, [2018](#)). It is based on Seligman's happiness theory (Seligman, [2002](#)) which addresses three main components of happiness: 1) pleasure or positive emotion, 2) engagement in activities, and 3) meaning in life. Considering Seligman et al. (2006)'s work, a six-session therapeutic plan was developed after slight modifications (techniques were chosen from PPT manual based on their relevancy to study participants and purpose instead of following the exact plan mentioned in Seligman et al. [2006](#)). This plan was approved by the original authors and aimed to enhance positive emotions, interpersonal relationships, engagement and meaning of life, rather than directly targeting presenting symptoms of psychopathology. The present research is designed to assess the effectiveness of six sessions of group positive psychotherapy intended to lessen depressive symptomatology by increasing pleasure, engagement, and focusing on meaning in life. In addition, it aimed at improving satisfaction with life among students.

Procedure

Before carrying out the study, it was first approved by the ethics committee of the Centre for Clinical Psychology, University of the Punjab. Following the approval, the study participants were approached online through advertisements using social media platforms due to the prevailing COVID situation. Those students who met the criteria and provided a written consent were included in the study. The participants were divided

in two groups that were matched on the basis of age, gender, education, and intensity of depression. Both groups were assessed at pre-intervention level using measures of depression and satisfaction with life. Later, the treatment group was given six sessions of positive psychotherapy in a group setting that included audio-recordings. This was carried in accordance with the guidelines provided in the PPT manual. However, the mode of intervention was online (with permission of the author), keeping in view the academic schedules of the students, their safety, and the concerns surrounding Coronavirus as well as the similar evidence from the literature (Yurayat & Seechaliao, [2021](#)).

The sessions were held twice a week, with each session consisting of 90 minutes. The sessions were carried out by a trainee clinical psychologist who was practicing traditional therapies. Although no specific training was required to administer the PPT protocol, to ensure the integrity of the process, the sessions were supervised by a consultant clinical psychologist who had more than ten years of experience in the field and manual adherence ratings by an independent field expert. To examine the efficacy of PPT in reducing depressive symptomatology and improving life satisfaction, both treatment and control groups were assessed at post-intervention level and their scores were compared. In addition, participants were asked to provide their subjective feedback on the four areas of well-being (individual, social, interpersonal, & general well-being) on a 0-10 scale following the PPT intervention.

The study data was analyzed using SPSS V23 by calculating descriptive statistics and through Mann-Whitney U and Wilcoxon signed rank tests.

Results

At first, the data were screened and then descriptive statistics and reliability values (i.e., Chronbach's alpha) were calculated for assessment measures and their dimensions.

Table 1

Descriptive Statistics and Reliability for Patient Health Questionnaire (PHQ-9) and Multidimensional Students' Life Satisfaction Scale (MSLSS) and its Dimensions (N=22)

Measures	<i>k</i>	<i>M</i>	<i>SD</i>	α	Skewness
PHQ-9	9	9.90	6.92	.86	0.581

Measures	<i>k</i>	<i>M</i>	<i>SD</i>	α	Skewness
MSLSS	40	183.77	27.63	.91	-0.586
Family	7	35.27	6.59	.87	-1.368
Friends	9	40.92	9.26	.85	-0.967
School	8	37.54	9.00	.90	-1.148
Environment	9	37.04	9.92	.85	-0.188
Self	7	33.00	6.95	.86	-1.066

As evident, Table 1 shows that both PHQ-9 and MSLSS are psychometrically reliable measures to assess depression and satisfaction with life.

Table 2

Mann-Whitney U Test Comparing Experimental and Control Group on Depression and Satisfaction with Life and its Domains at Pre-Intervention Level (N=22)

Scales	<i>M</i>	<i>SD</i>	Median	<i>U</i>	<i>p</i>
PHQ-9	9.90	2.52	10	58.0	.867
MSLSS	16.95	17.40	162	32.0	.061
Family	33.45	4.81	33.5	48.0	.410
Friends	36.95	8.20	36.5	51.0	.532
School	31.36	8.64	31.5	31.5	.056
Environment	32.63	6.97	51	51.0	.532
Self	30.40	5.88	45.5	45.5	.323

Note. PHQ-9=Patient Health Questionnaire, MSLSS= Multidimensional Student's Life Satisfaction Scale, *U*= Mann-Whitney U Test

The findings provided in Table 2 show that there was no significant difference between control and experimental groups when they were compared on the score of depression and satisfaction with life at pre-intervention level. This suggests that both groups were comparable.

Table 3

The Wilcoxon Signed Rank Test Comparing Experimental Group on Depression and Satisfaction with Life and its Domains at Pre and Post - Intervention Levels (N=11)

Scales	Pre-Intervention			Post-Intervention			<i>Z</i>	<i>p</i>	<i>d</i>
	<i>M</i>	<i>SD</i>	Median	<i>M</i>	<i>SD</i>	Median			
PHQ-9	9.90	2.73	11	3.0	2.14	3	-2.94	.003	.89
MSLSS	151.9	15.4	156	203	16.32	203	-2.94	.003	.89

Scales	Pre-Intervention			Post-Intervention			Z	p	d
	<i>M</i>	<i>SD</i>	Median	<i>M</i>	<i>SD</i>	Median			
Family	31.27	6.06	32	38.18	3.40	40	-2.81	.005	.85
Friends	35.90	4.18	36	43.81	6.17	43	-2.54	.011	.77
School	27.63	5.10	29	42.09	5.50	44	-2.95	.003	.89
Environment	32.90	4.15	32	40.63	6.60	41	-2.86	.004	.86
Self	28.90	2.87	30	38.27	2.86	40	-2.94	.003	.89

Note. PHQ-9=Patient Health Questionnaire; MSLSS= Multidimensional Student's Life Satisfaction Scale; Z= Wilcoxon signed rank.

Table 3 is showing the pre and post-assessment observations for the experimental group after 6-sessions of PPT. The findings showed that experiment group participants' scores differed in the post-intervention assessment. A significant decrease in depression was observed, along with an increased score in life satisfaction and its domains, including self, friends, living environment, and education.

Table 4

The Wilcoxon Signed Rank Test Comparing Control Group on Depression and Satisfaction with Life and its Domains at Pre and Post -Intervention Levels (N=11)

Scales	Pre-Intervention			Post-Intervention			Z	p
	<i>M</i>	<i>SD</i>	Median	<i>M</i>	<i>SD</i>	Median		
PHQ-9	9.90	2.42	10	11.5	3.55	10	-1.53	.126
MSLSS	170.3	19.72	169	167.8	17.22	168	-.178	.859
Family	34.27	4.26	35	36.54	3.50	37	-1.33	.182
Friends	38.0	11.02	37	37.27	9.42	38	-.153	.878
School	35.09	10.01	38	30.0	5.84	30	-1.37	.169
Environment	32.36	9.21	31	34.54	8.31	34	-.510	.610
Self	31.90	7.71	32	29.45	6.63	29	-1.02	.304

Note. PHQ-9=Patient Health Questionnaire; MSLSS= Multidimensional Student's Life Satisfaction Scale; Z= Wilcoxon Signed Rank

In comparison to the data presented in Table 3, the above Table 4 shows the pre and post-assessment scores for the control group, which was a no-treatment group. It was observed that the scores of participants in this group remained almost similar on both depression and satisfaction with life. There was no significant change in control group scores across the two points of observation. Overall, the findings suggested that the PPT proved to be effective in lowering the level of depression and enhancing satisfaction with life across various domains, including satisfaction with family, self, educational institute, friends and living environment.

Discussion

The present study was carried out with university students, and examined the utility of a PPT, which is considered a renowned therapy for cultivating positive emotions, engagement and meaning in life, and improving interpersonal relationships. The study investigated the efficacy of PPT in reducing the symptoms of depression and improving life satisfaction. While providing the evidence on the efficacy of PPT, the present study particularly focuses on the student population and highlights the need to address the mental health challenges that increasingly surface within academic organizations.

The findings of the study suggested that PPT can significantly reduce the symptoms of depression and lead to improvement in life satisfaction among the students. These findings are in alignment with the principles of PPT that emphasize positive emotions, strengths, and personal growth (Seligman, [2002](#); Seligman & Csikszentmihalyi, [2000](#)). Additionally, considering the strong emphasis of PPT on a client's strengths, growth, and well-being, it is convincing that PPT could be an efficacious and culturally sensitive approach, unlike other therapies focusing on symptoms of illness, for individuals living in collectivistic cultures like those in Asia. In such conventional cultures, harmony in relationships, integrity, and well-being of the group members are promoted (Shavitt & Zhang, [2004](#)), which is also the focus of PPT. Furthermore, the current study evidence is significantly supported by previous studies, which tested the efficacy of PPT with various pathological conditions, including depression (Asgharipoor et al., [2012](#); Goodwin, [2010](#); Mazhar & Riaz, [2020](#); Schrank et al., [2015](#); Uliaszek et al., [2016](#)). It highlights that PPT is an effective approach to reduce symptoms of psychopathology and improve satisfaction with life and well-being of individuals.

It has been tested previously that PPT significantly improves the positive psychological traits such as happiness and resilience, while promoting post-traumatic growth and reducing symptoms of depression (Woo et al., [2022](#)). This suggests that PPT not only reduces symptoms of pathology but also equips an individual with a set of traits and emotions that assist them in dealing with the aftermath of challenging events. Like in the present case, PPT has significantly reduced depression, and improved students' satisfaction with their lives. This is supported by sufficient evidence from the previous literature (Asgharipoor et al., [2012](#); Asl et al.,

[2016](#); Furchtlehner et al., [2024](#); Hoppen & Morina, [2021](#); Mazhar & Riaz, [2020](#); Seligman et al., [2006](#)). Depression is a reaction to adverse life events, such as experiencing a loss or failure (Freud, [1917](#); Jones et al., [2013](#)). In that, the present findings suggest that students receiving PPT could utilize their strengths in building confidence and resilience to win over their psychological distress resulting from MDCAT failure and deal with their current challenges, e.g., academic pressure, career related concerns, etc. This is because psychological factors like resilience buffer against the emotional turmoil experienced due to failures or mistakes (Johnson et al., [2017](#)). Another important factor which should be considered here is spiritual beliefs and coping, which have strong influence on improving the emotional well-being of the individuals (Rani & Ghazi, [2023](#); Salman et al., [2020](#)). It is likely that PPT remained efficacious in the current study because it strongly aligns with the spiritual beliefs and practices followed by majority of the people in Pakistan, for instance, gratitude and openminded attitude toward difficulties.

Evidence reports that mental illnesses such as depression are a prevalent issue in emerging economic countries, majorly due to low mental health literacy (Alvarez-Huerta et al., [2021](#)). The same goes for Pakistan as documented (Albarqi et al., [2024](#); Alvi et al., [2023](#)). In this context, the findings of this study call the attention of mental health stakeholders, policy makers, and the families to take necessary steps towards prevention of prevailing mental health difficulties among the student population, such as depression resulting from academic and family stressors. This is important because these stressors significantly influence their academic careers (Deng et al., [2022](#)).

The findings of this study also indicate PPT as a brief, cost-effective, and preventive approach that could be utilized with the Pakistani population, sometimes with slight modifications, to meet the specific population needs. Being a brief and strength-focused approach, PPT could be especially useful with the student population for accommodating their academic schedules and equally acknowledging their strengths and weaknesses in order to ensure their therapeutic engagement. Additionally, this perspective also encourages the use of PPT in sensitive situations where individuals are resistant and the chances of dropouts are higher.

Despite many strengths, the present study has a few limitations. First, this study was conducted with a smaller sample of students using the

English version of the protocol. So, it cannot be generalized that the translated version of the PPT protocol would be equally effective with other population groups as the original English version. In addition, the study didn't use randomization and has a potential for bias which further limits its generalizability. Furthermore, the researchers used a brief protocol and made efforts to ensure therapeutic alliance; however, the use of a digital mode of delivery might have influenced this process and, consequently, the study's findings. In the future, studies should focus on working with larger and different sample groups to test the efficacy of PPT. Moreover, comparative studies can be carried out to evaluate PPT in relation to conventional therapies that are currently practiced by clinicians, as well as different modes of therapeutic administration. This would provide valuable evidence on emerging therapies and help address gaps in the indigenous literature. The study also remained limited as researchers could not follow up with the participants. However, future studies could be planned with post therapy follow ups to track the efficacy of the PPT intervention.

Conclusion

In conclusion, the findings of the study suggested that PPT is an efficacious approach in managing depression and improving life satisfaction among the students who are experiencing mental health challenges due to emotionally debilitating experiences such as failures or trauma. The study contributes to the literature and encourages using PPT for its brevity and friendly approach.

Author Contribution

Amina Shahzadi: conceptualization, methodology, data curation, formal analysis, writing – original draft, writing – review & editing, visualization, project administration. **Faiza Safdar:** supervision, conceptualization, methodology.

Conflict of Interest

The authors of the manuscript have no financial or non-financial conflict of interest in the subject matter or materials discussed in this manuscript.

Data Availability Statement

The data associated with this study will be provided by the corresponding author upon request.

Funding Details

No funding has been received for this research.

Generative AI Disclosure Statement

The authors did not use any type of generative artificial intelligence software for this research.

References

Ahmad, S. S., & Koncsol, S. W. (2022). Cultural factors influencing mental health stigma: Perceptions of Mental Illness (POMI) in Pakistani emerging adults. *Religions*, 13(5), Article e401. <https://doi.org/10.3390/rel13050401>

Akhtar, P., Ma, L., Waqas, A., Naveed, S., Li, Y., Rahman, A., & Wang, Y. (2020). Prevalence of depression among university students in low and middle income countries (LMICs): A systematic review and meta-analysis. *Journal of Affective Disorders*, 274, 911–919. <https://doi.org/10.1016/j.jad.2020.03.183>

Albarqi, H., Aqeli, A. J., Alsaati, M. M., Alsaadawi, M., Basubrain, M. S., & Elessaw, L. R. (2024). A systematic review on inequalities in access to mental healthcare services in low and middle-income countries. *Rawal Medical Journal*, 49(1), Article e209.

Alvarez-Huerta, P., Muela, A., & Larrea, I. (2021). Student engagement and creative confidence beliefs in higher education. *Thinking Skills and Creativity*, 40(2), Article e100821. <http://dx.doi.org/10.1016/j.tsc.2021.100821>

Alvi, M. H., Ashraf, T., Naz, F., Sardar, A., Ullah, A., Patel, A., Kiran, T., Gumber, A., & Husain, N. (2023). Burden of mental disorders by gender in Pakistan: Analysis of global burden of disease study data for 1990–2019. *BJPsych Bulletin*, 48(6), 333–340. <https://doi.org/10.1192/bjb.2023.76>

Asgharipoor, N., Asgharnejad Farid, A., Arshadi, H., & Sahebi, A. (2012). A comparative study on the effectiveness of positive psychotherapy and group cognitive-behavioral therapy for the patients suffering from major depressive disorder. *Iranian Journal of Psychiatry and Behavioral Sciences*, 6(2), 33–41.

Asl, S. T. S., Sadeghi, K., Bakhtiari, M., Ahmadi, S. M., Nazari Anamagh, A., & Khayatan, T. (2016). Effect of group positive psychotherapy on improvement of life satisfaction and the quality of life in infertile woman. *International Journal of Fertility & Sterility*, 10(1), 105–112. <https://doi.org/10.22074/ijfs.2016.4775>

Barnett, P., Saunders, R., Buckman, J. E. J., Cardoso, A., Cirkovic, M., Leibowitz, J., Main, N., Naqvi, S. A., Singh, S., Stott, J., Varsani, L.,

Wheatly, J., & Pilling, S. (2022). Are students less likely to respond to routinely delivered psychological treatment? A retrospective cohort analysis. *Comprehensive Psychiatry*, 119, Article e152348. <https://doi.org/10.1016/j.comppsych.2022.152348>

Bilal, A., & Riaz, A. (2020). Academic stress and suicidal ideation in MDCAT repeating candidates: mediating role of depression and moderating role of age and gender. *Pakistan Journal of Physiology*, 16(3), 7–10.

Bukhari, S. R., Saba, F. (2017). Depression, anxiety and stress as negative predictors of life satisfaction in university students. *Rawal Medical Journal*, 42(2), 255–257.

Cook, S. C., Schwartz, A. C., & Kaslow, N. J. (2017). Evidence-based psychotherapy: Advantages and challenges. *Neurotherapeutics: The Journal of the American Society for Experimental NeuroTherapeutics*, 14(3), 537–545. <https://doi.org/10.1007/s13311-017-0549-4>

Cuijpers, P., Cristea, I. A., Ebert, D. D., Koot, H. M., Auerbach, R. P., Bruffaerts, R., & Kessler, R. C. (2016). Psychological treatment of depression in college students: A metaanalysis. *Depression and Anxiety*, 33(5), 400–414. <https://doi.org/10.1002/da.22461>

Deng, Y., Cherian, J., Khan, N. U. N., Kumari, K., Sial, M. S., Comite, U., Gavurova, B., & Popp, J. (2022). Family and academic stress and their impact on students' depression level and academic performance. *Frontiers in Psychiatry*, 13, Article e869337. <https://doi.org/10.3389/fpsyg.2022.869337>

Dowlatabadi, M. M., Ahmadi, S. M., Sorbi, M. H., Beiki, O., Razavi, T. K., & Bidaki, R. (2016). The effectiveness of group positive psychotherapy on depression and happiness in breast cancer patients: A randomized controlled trial. *Electronic Physician*, 8(3), 2175–2180. <https://doi.org/10.19082/2175>

Erschens, R., Keifenheim, K. E., Herrmann-Werner, A., Loda, T., Schwille-Kiuntke, J., Bugaj, T. J., Nikendei, C., Huhn, D., Zipfel, S., & Junne, F. (2018). Professional burnout among medical students: Systematic literature review and meta-analysis. *Medical Teacher*, 41(2), 172–183. <https://doi.org/10.1080/0142159X.2018.1457213>

Fatima, I., Nadeem, M., Naeem, M. Q., Raza, H. M. Z. (2021). Impact of MDCAT scores on pessimism, loneliness, and depressive symptoms: Moderating role of coping strategies. *World Journal of Advanced Research and Reviews*, 12(2), 354–364. <https://doi.org/10.30574/wjarr.2021.12.2.0600>

Feng, T., Jia, X., Pappas, L., Zheng, X., Shao, T., Sun, L., Weisberg, C., Li, M. L., Rozelle, S., & Ma, Y. (2022). Academic performance and the link with depressive symptoms among Rural Han and Minority Chinese adolescents. *International Journal of Environmental Research and Public Health*, 19(10), Article e6026. <https://doi.org/10.3390/ijerph19106026>

Ford, B. Q., Dmitrieva, J. O., Heller, D., Chentsova-Dutton, Y., Grossmann, I., Tamir, M., Uchida, Y., Koopmann-Holm, B., Floerke, V. A., Uhrig, M., Bokhan, T., & Mauss, I. B. (2015). Culture shapes whether the pursuit of happiness predicts higher or lower well-being. *Journal of Experimental Psychology. General*, 144(6), 1053–1062. <https://doi.org/10.1037/xge0000108>

Fredrickson B. L. (2004). The broaden-and-build theory of positive emotions. *Philosophical transactions of the Royal Society of London. Series B, Biological Sciences*, 359(1449), 1367–1378. <https://doi.org/10.1098/rstb.2004.1512>

Freud, S. (1917). Mourning and melancholia. In J. Stratchey (trans.), *The standard edition of the complete psychological works of Sigmund Freud* (Volume 14, pp. 237–258). Hogarth.

Furchtlehner, L.M., Fischer, E., Schuster, R., & Laireiter, A-R. (2024). A comparative study on the efficacy of group positive psychotherapy and group cognitive behavioral therapy on flourishing, happiness and satisfaction with life: A randomized controlled trial. *Journal of Happiness Studies*, 25, Article e104. <https://doi.org/10.1007/s10902-024-00806-y>

Goodwin, E. M. (2010). *Does group positive psychotherapy help improve relationship satisfaction in a stressed and/or anxious population?* (Publication No. 3428275) [Doctoral dissertation, Palo Alto University]. ProQuest Dissertations and Theses.

Hoppen, T. H., & Morina, N. (2021). Efficacy of positive psychotherapy in reducing negative and enhancing positive psychological outcomes: A meta-analysis of randomised controlled trials. *BMJ Open*, 11(9), e046017. <https://doi.org/10.1136/bmjopen-2020-046017>

Huebner, E.S. (2001). *Manual for the multidimensional students' life satisfaction scale*. University of South Carolina.

Irie, T., & Yokomitsu, K. (2019). Relationship between dispositional mindfulness and living condition and the well-being of first-year university students in Japan. *Frontier in Psychology*, 10, Article e489191. <https://doi.org/10.3389/fpsyg.2019.02831>

Johnson, J., Panagioti, M., Bass, J., Ramsey, L., & Harrison, R. (2017). Resilience to emotional distress in response to failure, error, or mistakes: A systematic review. *Clinical Psychology Review*, 52, 19–42. <https://doi.org/10.1016/j.cpr.2016.11.007>

Jones, N. P., Papadakis, A. A., Orr, C. A., & Strauman, T. J. (2013). Cognitive processes in response to goal failure: A study of ruminative thought and its affective consequences. *Journal of Social and Clinical Psychology*, 32(5), 482–503. <https://doi.org/10.1521/jscp.2013.32.5.482>

Kabir, R., Syed, H. Z., Vinnakota, D., Okello, S., Isigi, S. S., Abdul Kareem, S. K., Parsa, A. D., & Arafat, S. M. Y. (2024). Suicidal behaviour among the university students in the UK: A systematic review. *Helijon*, 10(2), Article e24069 <https://doi.org/10.1016/j.heliyon.2024.e24069>

Kahler, C. W., Spillane, N. S., Day, A., Clerkin, E. M., Parks, A., Leventhal, A. M., et al. (2014). Positive psychotherapy for smoking cessation: Treatment development, feasibility, and preliminary results. *Journal of Positive Psychology*, 9(1), 19–29. <https://doi.org/10.1080/17439760.2013.826716>

Khan, M. N., Akhtar, P., Ijaz, S., & Waqas, A. (2021). Prevalence of depressive symptoms among university students in Pakistan: A systematic review and meta-analysis. *Frontiers in Public Health*, 8, Article e603357. <https://doi.org/10.3389/fpubh.2020.603357>

Kim, Y. E., & Lee, B. (2019). The psychometric properties of the Patient Health Questionnaire-9 in a sample of Korean university students.

Psychiatry Investigation, 16(12), 904–910.
<https://doi.org/10.30773/pi.2019.0226>

Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>

Liu, S. S., Shteynberg, G., Morris, M. W., Yang, Q., & Galinsky, A. D. (2021). How does collectivism affect social interactions? A test of two competing accounts. *Personality & Social Psychology Bulletin*, 47(3), 362–376. <https://doi.org/10.1177/0146167220923230>

Liu, X., & Wang, J. (2024). Depression, anxiety, and student satisfaction with university life among college students: A cross-lagged study. *Humanities and Social Sciences Communications*, 11, e1172. <https://doi.org/10.1057/s41599-024-03686-y>

Lukaschek, K., Vanajan, A., Johar, H., Weiland, N., & Ladwig, K. H. (2017). "In the mood for ageing": Determinants of subjective well-being in older men and women of the population-based KORA-Age study. *BMC Geriatrics*, 17(1), Article e126. <https://doi.org/10.1186/s12877-017-0513-5>

Mathew, N., Khakha, D. C., Qureshi, A., Sagar, R., & Khakha, C. C. (2015). Stress and coping among adolescents in selected schools in the capital city of India. *Indian Journal of Pediatrics*, 82(9), 809–816. <https://doi.org/10.1007/s12098-015-1710-x>

Mazhar, A. F., & Riaz, M. N. (2020). Effectiveness of positive psychotherapy for young adults with depressive symptoms. *The Journal of the Pakistan Medical Association*, 70(5), 856–859. <https://doi.org/10.5455/JPMA.16572>

Memon, R., Asif, M., Khoso, A. B., Tofique, S., Kiran, T., Chaudhry, N., Husain, N., Edwards, S. J. L., & Pakistan Institute of Living and Learning. (2021). Recognising values and engaging communities across cultures: Towards developing a cultural protocol for researchers. *BMC Medical Ethics*, 22(1), Article e47. <https://doi.org/10.1186/s12910-021-00608-4>

Mofatteh, M. (2020). Risk factors associated with stress, anxiety, and depression among university undergraduate students. *AIMS Public Health*, 8(1), 36–65. <https://doi.org/10.3934/publichealth.2021004>

Moir, F., Yielder, J., Sanson, J., & Chen, Y. (2018). Depression in medical students: Current insights. *Advances in Medical Education and Practice*, 9, 323–333. <https://doi.org/10.2147/AMEP.S137384>

Muneer, A., Riaz, M., Masood, I., Akbar, J., Ahmad, S., Hussain, T., Sehar, G., Gillani, A. H., & Khan, Z. (2024). Incidence of depression, anxiety, and stress among students of health sciences and non-health sciences: A comparative cross-sectional study from Pakistan. *International Journal of Pharmacy & Integrated Health Sciences*, 6(1), 25–36. <https://doi.org/10.56536/ijpihs.v6i1.170>

National Academies of Sciences, Engineering, and Medicine. (2019). *The promise of adolescence: Realizing opportunity for all youth*. The National Academies Press.

Ooi, P. B., Khor, K. S., Tan, C. C., & Ong, D. L. T. (2022). Depression, anxiety, stress, and satisfaction with life: Moderating role of interpersonal needs among university students. *Frontiers in Public Health*, 10, Article e958884. <https://doi.org/10.3389/fpubh.2022.958884>

Rahman, R., Sheehan, D., Javed, A., Ahmad, S. H., Shafiq, K., Kanwal, U., Afridi, M. I., Nizami, A. T., Rasool, G., Taj, R., Ansari, M., Naim, M., Farooq, S., Memon, A., Karim, M. S., Hana, Y., & Ayub, M. (2024). *The National Psychiatric Morbidity Survey of Pakistan (2022): Prevalence, sociodemographic and disability correlates*. medRxiv. <https://doi.org/10.1101/2024.07.08.24310056>

Rani, A., & Ghazi, M. H. (2023). Relationship between spirituality and mental health conditions among undergraduate students. *Pakistan Journal of Psychological Research*, 38(4), 675– 687. <https://doi.org/10.33824/PJPR.2023.38.4.39>

Rashid, T. & Seligman, M. (2018). *Positive psychotherapy: Clinical manual*. Oxford University Press.

Rashid, T. (2014). Positive psychotherapy: A strength-based approach. *The Journal of Positive Psychology*, 10(1), 25–40. <https://doi.org/10.1080/17439760.2014.920411>

Rathod, S., Pinninti, N., Irfan, M., Gorczynski, P., Rathod, P., Gega, L., & Naeem, F. (2017). Mental health service provision in low- and middle-income countries. *Health Services Insights*, 10, 1–7. <https://doi.org/10.1177/1178632917694350>

Saldivia, S., Aslan, J., Cova, F., Bustos, C., Inostroza, C., & Castillo-Carreño, A. (2023). Life satisfaction, positive affect, depression and anxiety symptoms, and their relationship with sociodemographic, psychosocial, and clinical variables in a general elderly population sample from Chile. *Frontiers in Psychiatry*, 14, Article e1203590. <https://doi.org/10.3389/fpsy.2023.1203590>

Salman, M., Asif, N., Mustafa, Z. U., Khan, T. M., Shehzadi, N., Tahir, H., Raza, M. H., Khan, M. T., Hussain, K., Khan, Y. H., Butt, M. H., & Mallhi, T. H. (2020). Psychological impairment and coping strategies during the COVID-19 pandemic among students in Pakistan: A cross-sectional analysis. *Disaster Medicine and Public Health Preparedness*, 16(3), 920–926. <https://doi.org/10.1017/dmp.2020.397>

Schnettler, B., Orellana, L., Sepúlveda, J., Miranda, H., Grunert, KG, Lobos, G., & Hueche, C. (2017). Psychometric properties of the Multidimensional Students' Life Satisfaction Scale in a sample of Chilean university students. *Suma Psicológica*, 24, 97–106. <https://doi.org/10.1016/j.sumpsi.2017.06.001>

Schrank, B., Brownell, T., Jakaite, Z., Larkin, C., Pesola, F., Riches, S., et al. (2015). Evaluation of a positive psychotherapy group intervention for people with psychosis: Pilot randomized controlled trial. *Epidemiology and Psychiatric Sciences*, 25(3), 235–246. <https://doi.org/10.1017/S2045796015000141>

Seligman, M. E. P. (2002). *Authentic happiness: Using the new positive psychology to realize your potential for lasting fulfillment*. Free Press.

Seligman, M. E. P. (2011). *Flourish: A visionary new understanding of happiness and well-being*. Free Press.

Seligman, M. E. P., Rashid, T., & Parks, A. C. (2006). Positive psychotherapy. *The American Psychologist*, 61(8), 774–788. <https://doi.org/10.1037/0003-066X.61.8.774>

Seligman, M. E., & Csikszentmihalyi, M. (2000). Positive psychology. An introduction. *The American Psychologist*, 55(1), 5–14. <https://doi.org/10.1037/0003-066x.55.1.5>

Shavitt, S., & Zhang, J. (2004). Advertising and culture. In Charles D. S. (Ed.), *Encyclopedia of applied psychology* (pp. 47–51). Elsevier.

Silva, S. A., Silva, S. U., Ronca, D. B., Gonçalves, V. S. S., Dutra, E. S., & Carvalho, K. M. B. (2020). Common mental disorders prevalence in adolescents: A systematic review and meta-analyses. *PLOS ONE*, 15(4), Article e0232007. <https://doi.org/10.1371/journal.pone.0232007>

Sinval, J., Oliveira, P., Novais, F., Almeida, C. M., & Telles-Correia, D. (2024). Exploring the impact of depression, anxiety, stress, academic engagement, and dropout intention on medical students' academic performance: A prospective study. *Journal of Affective Disorders*, 368, 665–673. <https://doi.org/10.1016/j.jad.2024.09.116>

Storrie, K., Ahern, K., & Tuckett, A. (2010). A systematic review: Students with mental health problems--a growing problem. *International Journal of Nursing Practice*, 16(1), 1–6. <https://doi.org/10.1111/j.1440-172X.2009.01813.x>

Substance Abuse and Mental Health Services Administration. (2022). *Key substance use and mental health indicators in the United States: Results from the 2021 National Survey on Drug Use and Health* (HHS Publication No. PEP22-07-01-005, NSDUH Series H-57). <https://www.samhsa.gov/data/report/2021-nsduh-annual-national-report>

Tam, W., Lo, K., & Pacheco, J. (2019). Prevalence of depressive symptoms among medical students: Overview of systematic reviews. *Medical Education*, 53(4), 345–354. <https://doi.org/10.1111/medu.13770>

Uliaszek, A. A., Rashid, T., Williams, G. E., & Gulamani, T. (2016). Group therapy for university students: A randomized control trial of dialectical behavior therapy and positive psychotherapy. *Behaviour Research and Therapy*, 77, 78–85. <https://doi.org/10.1016/j.brat.2015.12.003>

Ullah, I., Ali, S., Ashraf, F., Hakim, Y., Ali, I., Ullah, A. R., Chattu, V. J., & Pakpour, A. H. (2022). Prevalence of depression and anxiety among general population in Pakistan during COVID-19 lockdown: An online-

survey. *Current Psychology*, 43, 8338–8345.
<https://doi.org/10.1007/s12144-022-02815-7>

Walsh, S., Cassidy, M., & Priebe, S. (2017). The application of positive psychotherapy in mental health care: A systematic review. *Journal of Clinical Psychology*, 73(6), 638–651.
<https://doi.org/10.1002/jclp.22368>

Woo, M-S., Woo, J-H., & Yang, H-C. (2022). The effect of positive psychotherapy (PPT) programs on participants' happiness and resilience. *Journal of Economics Marketing, and Management*, 10(5), 15–24. <http://dx.doi.org/10.20482/jemm.2022.10.5.15>

Yousef, S., Athamneh, M., Masuadi, E., Ahmad, H., Loney, T., Moselhy, H. F., Al-Maskari, F., & ElBarazi, I. (2017). Association between depression and factors affecting career choice among Jordanian nursing students. *Frontiers in Public Health*, 5, Article e311.
<https://doi.org/10.3389/fpubh.2017.00311>

Yurayat, P., & Seechaliao, T. (2021). Effectiveness of online positive psychology intervention on psychological well-being among undergraduate students. *Journal of Education and Learning*, 10(4), 143–155. <https://doi.org/10.5539/jel.v10n4p143>