



Scientific Inquiry and Review (SIR)

Volume 2, Issue 2, April 2018

ISSN (P): 2521-2427, ISSN (E): 2521-2435

Journal DOI: <https://doi.org/10.29145/sir>

Issue DOI: <https://doi.org/10.29145/sir/22>

Homepage: <https://ssc.umt.edu.pk/sir/Home.aspx>

Journal QR Code:



Article:

Modified Dust-Lower-Hybrid Waves in Quantum Plasma

Author(s):

Abdul Rauf
Israr Zeb
Muhammad Saqlain

Online Published:

April 2018

Article DOI:

doi.org/10.29145/sir/22/020202

Article QR Code:



Abdul Rauf

To cite this article:

Rauf A, Zeb I, Saqlain M. Modified dust-lower-hybrid waves in quantum plasma. *Sci Inquiry Rev.* 2018;2(2):12–21. DOI: <https://doi.org/10.29145/sir/22/020202>



A publication of the
School of Science
University of Management and Technology
Lahore

Modified Dust-Lower-Hybrid Waves In Quantum Plasma

Abdul Rauf ¹, I. Zeba ², Muhammad Saqlain ³

¹ Department of Physics, LGU, Lahore, Pakistan

² Department of Physics, LCW University Lahore-54000, Pakistan

³ Department of Mathematics, Lahore Garrison University, Lahore, Pakistan

Abstract

Dust-lower-hybrid waves in quantum plasma have been studied. The dispersion relation of the dust-lower-hybrid wave has been examined using the quantum hydrodynamic model of plasma in an ultra-cold Fermi dusty plasma in the presence of a uniform external magnetic field. Graphical analysis shows that the electron Fermi temperature effect and the quantum corrections give rise to significant effects on the dust-lower-hybrid wave of the magnetized quantum dusty plasma.

Keywords: Magneto plasma, wavelength, microelectronics, hydrodynamics, quantum mechanics

1. Introduction

Quantum mechanical effects in some specific areas of plasma physics have great significance. When plasma is cooled to an extremely low temperature, the de Broglie wavelengths of the plasma particles are comparable to the dimensions of the system. In such plasmas, the ultra-cold dense plasma would behave as a Fermi gas and quantum mechanical effects might play a vital role in the behaviour of the charge carriers of plasmas under these conditions. In microelectronics [1,2] and very large integrated circuit fabrications, the system may develop contaminants due to etching, implantations etc. which may lead to new properties. The laser produced plasmas [3] and plasmas in high density astrophysical objects [4-6] may also be contaminated by various reasons. Thus, these ultra-cold plasma systems may behave as dusty plasmas where quantum mechanical effects could unveil the properties of these systems. This research aims to present the quantum effects on dust-lower-hybrid waves, where dust dynamic plays a vital role. Within the ambit of a symmetrical external magnetic field, the modification of the dust-lower-hybrid wave in quantum plasma can be achieved by following the standard techniques [7-10].

2. Material and Method

The equations in the quantum hydrodynamic model for electrons and ions of dusty plasma in presence of a dust-lower-hybrid perturbation and \vec{B}_0 are

$$\frac{\partial n_{j1}}{\partial t} + n_{j0}(\vec{\nabla} \cdot \vec{V}_{j1}) = 0 \quad (1)$$

n_{j1} = perturb particle density of jth species

n_{j0} = equilibrium particle density of jth species

\vec{V}_{j1} = perturb velocity of jth species

The equation of motion for electron and ions of dusty plasma in the presence of perturbation and \vec{B}_0 is

$$m_j n_{j0} \frac{\partial \vec{v}_{j1}}{\partial t} = n_{j0} q_j (\vec{E}_1 + \vec{v}_{j1} \times \vec{B}_0) \quad (2)$$

m_j = mass of jth species

q_j = charge of jth species

\vec{E}_1 = electric field intensity in the presence of a DLH perturbation

\vec{B}_0 = equilibrium/uniform magnetic field.

$$\vec{E}_1 = -\vec{\nabla} \phi_1$$

ϕ_1 = perturb electrostatic potential

and

$$\frac{q_j \vec{B}_0}{m_j} = \vec{\omega}_{cj}$$

So equation (2) becomes

$$\frac{\partial}{\partial t} \vec{v}_{j1} = -\frac{q_j}{m_j} \vec{\nabla} \phi_1 + \vec{v}_{j1} \times \vec{\omega}_{cj} \quad (3)$$

Dielectric susceptibility of plasma can be obtained by solving the equations (1) and (3)

according to Fourier Laplace transformation

$$\begin{aligned} \frac{\partial}{\partial t} &= -i\omega \quad \text{and} \quad \vec{\nabla} = i\vec{k} \\ \vec{v}_{j1} &= \vec{v}_\perp + \vec{v}_\parallel \\ \vec{k} &= \vec{k}_\perp + \vec{k}_\parallel \end{aligned}$$

From equation (3)

$$\vec{v}_\perp = \frac{i q_j \phi_1 [(\vec{k}_\perp \times \vec{\omega}_{cj}) - i \vec{k}_\perp \omega]}{m_j (\omega^2 - \omega_{cj}^2)}$$

And

$$\vec{v}_\parallel = \frac{q_j \phi_1 \vec{k}_\parallel}{\omega m_j}$$

From equation (1)

$$\begin{aligned} n_{j1} &= \frac{n_{j0}}{\omega} [\vec{k}_\perp \cdot \vec{v}_\perp + \vec{k}_\parallel \cdot \vec{v}_\parallel] \\ n_{j1} &= \frac{n_{j0}}{\omega} \left[\frac{k_\perp^2 \omega q_j \phi_1}{m_j (\omega^2 - \omega_{cj}^2)} + \frac{k_\parallel^2 q_j \phi_1}{\omega m_j} \right] \quad (4) \end{aligned}$$

As

$$n_{j1} = -\frac{\chi_j k^2 \phi_1}{4\pi q_j} \quad (5)$$

From equations (4) and (5)

$$\chi_j = \frac{k_\perp^2}{k^2} \times \frac{4\pi n_{j0} q_j^2}{m_j (\omega^2 - \omega_{cj}^2)} - \frac{k_\parallel^2}{k^2} \times \frac{4\pi n_{j0} q_j^2}{\omega^2 m_j}$$

Since

$$\omega_{pj}^2 = \frac{4\pi n_{j0} q_j^2}{m_j} \quad (6)$$

Therefore

$$\chi_j = \frac{k_{\perp}^2}{k^2} \times \frac{\omega_{pj}^2}{(\omega_{cj}^2 - \omega^2)} - \frac{k_{\parallel}^2}{k^2} \times \frac{\omega_{pj}^2}{\omega^2} \quad (7)$$

For un-magnetized dust

$$\omega_{cd} = 0$$

$$\chi_d = -\frac{k_{\perp}^2}{k^2} \times \frac{\omega_{pd}^2}{\omega^2} - \frac{k_{\parallel}^2}{k^2} \times \frac{\omega_{pd}^2}{\omega^2}$$

So, the susceptibility for the unmagnetized and cold dust particle is

$$\chi_d = -\frac{\omega_{pd}^2}{\omega^2} \quad (8)$$

The dispersion relation is

$$\varepsilon(\omega, \vec{k}) = \varepsilon = 1 + \chi_e + \chi_i + \chi_d = 0 \quad (9)$$

$$1 + \left(\frac{k_{\perp}^2}{k^2} \times \frac{\omega_{pe}^2}{(\omega_{ce}^2 - \omega^2)} - \frac{k_{\parallel}^2}{k^2} \times \frac{\omega_{pe}^2}{\omega^2} \right)$$

$$+ \left(\frac{k_{\perp}^2}{k^2} \times \frac{\omega_{pi}^2}{(\omega_{ci}^2 - \omega^2)} - \frac{k_{\parallel}^2}{k^2} \times \frac{\omega_{pi}^2}{\omega^2} \right) + \left(-\frac{\omega_{pd}^2}{\omega^2} \right) = 0$$

$$\omega_{ce}^2 \gg \omega^2 \text{ and } \omega_{ci}^2 \gg \omega^2$$

Also

$$\omega_{pe} \gg \omega_{pi}$$

$$\omega_{pi}^2 \approx 0$$

$$\omega_{pi}^2 \gg \omega_{ci}^2$$

$$\omega^2 = \frac{k_{\perp}^2}{k^2} \times \frac{\omega_{pd}^2 \omega_{ci}^2}{\omega_{pi}^2} \left(1 + \frac{k_{\parallel}^2 \omega_{pe}^2}{k^2 \omega_{pd}^2} \right)$$

As, $k^2 = k_{\perp}^2 + k_{\parallel}^2$ if $k_{\perp}^2 \gg k_{\parallel}^2$ then, $k^2 = k_{\perp}^2$ or $\frac{k_{\parallel}^2}{k_{\perp}^2} = 1$

$$\omega^2 = \frac{\omega_{pd}^2 \omega_{ci}^2}{\omega_{pi}^2} \left(1 + \frac{k_{\parallel}^2 \omega_{pe}^2}{k^2 \omega_{pd}^2} \right)$$

$$\omega^2 = \omega_{dlh}^2 \left(1 + \frac{k_{\parallel}^2 \omega_{pe}^2}{k^2 \omega_{pd}^2} \right) \quad (10)$$

is the dispersion relation for DLH wave.

Now, equation (2) with quantum effect becomes

$$m_j n_{j0} \frac{\partial \vec{V}_{j1}}{\partial t} = n_{j0} q_j (\vec{E}_1 + \vec{V}_{j1} \times \vec{B}_0) - \vec{\nabla} P_{j1} + \frac{\hbar^2 \vec{\nabla}(\nabla^2 n_{j1})}{4 m_j} \quad (11)$$

$\frac{\hbar^2 \vec{\nabla}(\nabla^2 n_{j1})}{m_j}$ = Bohm Potential term

$\vec{\nabla} P_{j1}$ = Fermi pressure term

$$\frac{\partial \vec{V}_{j1}}{\partial t} = \frac{q_j}{m_j} (-\vec{\nabla} \phi_1 + \vec{V}_{j1} \times \vec{B}_0) - \frac{\vec{\nabla} P_{j1}}{m_j n_{j0}} + \frac{\hbar^2 \vec{\nabla}(\nabla^2 n_{j1})}{4 m_j^2 n_{j0}}$$

where

$$P_{j1} = 2 n_{j1} k_B T_{Fj}$$

▪ by assuming that electron and ions possess significant quantum mechanical effects

- by neglecting the quantum effects on dusty particles (because of high mass which gives to an insignificant de Broglie wavelength).

Since

$$\begin{aligned} \frac{\partial}{\partial t} &= -i\omega \quad \text{and} \quad \vec{\nabla} = i\vec{k}; \quad \nabla^2 = (i\vec{k})^2 = -k^2 \\ -i\omega \vec{V}_{j1} &= -\frac{q_j}{m_j} i\vec{k} \varphi_1 + \frac{q_j}{m_j} (\vec{V}_{j1} \times \vec{B}_o) - \frac{i\vec{k} (2 n_{j1} k_B T_{Fj})}{m_j n_{jo}} \\ &\quad + \frac{\hbar^2 (i\vec{k}) (-k^2 n_{j1})}{4 m_j^2 n_{jo}} \\ \omega \vec{V}_{j1} &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i \frac{q_j}{m_j} (\vec{V}_{j1} \times \vec{B}_o) + \frac{2\vec{k}}{n_{jo}} \left(\frac{k_B T_{Fj}}{m_j} \right) n_{j1} + \frac{\hbar^2 (\vec{k}) (k^2)}{4 m_j^2 n_{jo}} n_{j1} \\ \omega \vec{V}_{j1} &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i \frac{q_j}{m_j} (\vec{V}_{j1} \times \vec{B}_o) + \frac{\vec{k} V_{Fj}^2}{n_{jo}} (1 + \gamma_j) n_{j1} \end{aligned}$$

where

$$\begin{aligned} V_{Fj} &= \left(\frac{2 k_B T_{Fj}}{m_j} \right)^{\frac{1}{2}} \quad \text{and} \quad \gamma_j \\ &= \frac{\hbar^2 (k^2)}{8 m_j k_B T_{Fj}} \quad (\text{Quantum correction}) \\ \omega \vec{V}_{j1} &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i (\vec{V}_{j1} \times \vec{\omega}_{cj}) + \frac{\vec{k} V_{Fj}^2}{n_{jo}} n_{j1} \quad (12) \end{aligned}$$

where

$$V_{Fj}'^2 = V_{Fj}^2 (1 + \gamma_j)$$

From equation (1)

$$\begin{aligned} -i\omega n_{j1} + n_{jo} (i\vec{k} \cdot \vec{V}_{j1}) &= 0 \\ -\omega n_{j1} + n_{jo} (\vec{k} \cdot \vec{V}_{j1}) &= 0 \\ n_{j1} &= \frac{n_{jo}}{\omega} \vec{k} \cdot \vec{V}_{j1} \quad (13) \end{aligned}$$

So equation (12) becomes

$$\begin{aligned} \omega \vec{V}_{j1} &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i (\vec{V}_{j1} \times \vec{\omega}_{cj}) + \frac{\vec{k} V_{Fj}'^2}{n_{jo}} \left(\frac{n_{jo}}{\omega} \vec{k} \cdot \vec{V}_{j1} \right) \\ \omega \vec{V}_{j1} &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i (\vec{V}_{j1} \times \vec{\omega}_{cj}) + \frac{k^2 V_{Fj}'^2}{\omega} (\vec{V}_{j1}) \\ \omega \left(1 - \frac{k^2 V_{Fj}'^2}{\omega^2} \right) \vec{V}_{j1} &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i (\vec{V}_{j1} \times \vec{\omega}_{cj}) \end{aligned}$$

where

$$\begin{aligned} F'_j &= 1 - \frac{k^2 V_{Fj}'^2}{\omega^2} \\ \omega F'_j (\vec{V}_{j1}) &= \frac{q_j}{m_j} \vec{k} \varphi_1 + i (\vec{V}_{j1} \times \vec{\omega}_{cj}) \quad (14) \end{aligned}$$

Taking perpendicular (\perp) component of equation (14)

$$\omega F'_j (\vec{V}_{j\perp}) = \frac{q_j}{m_j} \vec{k}_\perp \phi_1 + i (\vec{V}_{j\perp} \times \vec{\omega}_{cj}) \quad (15)$$

Multiply this equation by $\vec{\omega}_{cj}$, we get

$$\begin{aligned} \omega F'_j (\vec{V}_{j\perp} \times \vec{\omega}_{cj}) &= \frac{q_j \phi_1}{m_j} (\vec{k}_\perp \times \vec{\omega}_{cj}) + i (\vec{V}_{j\perp} \times \vec{\omega}_{cj}) \times \vec{\omega}_{cj} \\ -i \omega F'_j \left(-i \omega F'_j \vec{V}_{j\perp} + \frac{i q_j \vec{k}_\perp \phi_1}{m_j} \right) &= -i \frac{q_j \phi_1}{m_j} (\vec{k}_\perp \times \vec{\omega}_{cj}) + \\ (\vec{V}_{j\perp} \times \vec{\omega}_{cj}) \times \vec{\omega}_{cj} \quad (16) \end{aligned}$$

Since

$$(\vec{V}_{j\perp} \times \vec{\omega}_{cj}) \times \vec{\omega}_{cj} = -\omega_{cj}^2 \vec{V}_{j\perp}$$

So last equation becomes

$$\vec{V}_{j\perp} = \frac{i q_j \phi_1 [\vec{k}_\perp \times \vec{\omega}_{cj} - i \omega F'_j \vec{k}_\perp]}{m_j [\omega^2 F_j'^2 - \omega_{cj}^2]} \quad (17)$$

Taking parallel (\parallel) component of equation (14), and putting $\vec{V}_{j\parallel} \times \vec{\omega}_{cj} = 0$

$$\omega F'_j \vec{V}_{j\parallel} = \frac{q_j \vec{k}_\parallel \phi_1}{m_j} + 0$$

$$\vec{V}_{j\parallel} = \frac{q_j \vec{k}_\parallel \phi_1}{\omega F'_j m_j} \quad (18)$$

From equation (1),

$$\begin{aligned} \frac{\partial n_{j1}}{\partial t} &= -n_{jo} (\vec{\nabla} \cdot \vec{V}_{j1}) \\ -i \omega n_{j1} &= -n_{jo} i (\vec{k}_\parallel + \vec{k}_\perp) (\vec{V}_{j\parallel} \cdot \vec{V}_{j\perp}) \\ n_{j1} &= \frac{n_{jo}}{\omega} (\vec{k}_\parallel \cdot \vec{V}_{j\parallel} + \vec{k}_\perp \cdot \vec{V}_{j\perp}) \end{aligned}$$

Putting the values of $\vec{V}_{j\perp}$ and $\vec{V}_{j\parallel}$, we get

$$\chi_j = \frac{k_\perp^2}{k^2} \frac{\omega_{pj}^2 F'_j}{(\omega_{cj}^2 - \omega^2 F_j'^2)} - \frac{k_\parallel^2}{k^2} \frac{\omega_{pj}^2}{F'_j \omega^2} \quad (19)$$

Let us consider a super-cooled magnetized Fermi dusty plasma where

- Electrons are considered hot at Fermi temperature and quantum
- Ions are cold and non-quantum
- Dust particles are cold and non-quantum

Then

$$\begin{aligned} \chi_e &= \frac{k_\perp^2}{k^2} \frac{\omega_{pe}^2 F'_e}{(-\omega^2 F_e'^2)} - \frac{k_\parallel^2}{k^2} \frac{\omega_{pe}^2}{F'_e \omega^2} \\ &= -\frac{k_\perp^2}{k^2} \frac{\omega_{pe}^2}{F'_e} - \frac{k_\parallel^2}{k^2} \frac{\omega_{pe}^2}{F'_e \omega^2} \\ &= -\frac{(k_\perp^2 + k_\parallel^2)}{k^2} \frac{\omega_{pe}^2}{F'_e \omega^2} \end{aligned}$$

Since $k_{\perp}^2 + k_{\parallel}^2 = k^2$

$$\chi_e = - \frac{\omega_{pe}^2}{F'_e \omega^2}$$

Since $F'_e = 1 - \frac{k^2 V_{Fe}^2}{\omega^2}$

$$\chi_e = - \frac{\omega_{pe}^2}{\omega^2 \left(1 - \frac{k^2 V_{Fe}^2}{\omega^2} \right)}$$

Assume $\omega^2 \ll k^2 V_{Fe}^2$, and $F'_e = - \frac{k^2 V_{Fe}^2}{\omega^2}$ then,

$$\begin{aligned} \chi_e &= - \frac{\omega_{pe}^2}{\omega^2 \left(- \frac{k^2 V_{Fe}^2}{\omega^2} \right)} \\ &= \frac{\omega_{pe}^2}{k^2 V_{Fe}^2} \end{aligned}$$

Since $\lambda_{Fe}^2 = \frac{V_{Fe}^2}{\omega_{pe}^2}$

λ'_{Fe} = Debye length of electrons at Fermi temperature

V'_{Fe} = Fermi speed of electrons

ω_{pe} = electron plasma frequency

So,
$$\chi_e = \frac{1}{k^2 \lambda_{Fe}^2} \quad (20)$$

Now for χ_i , $F'_i = 1$

$$\chi_i = \frac{K_{\perp}^2 \omega_{pi}^2}{K^2 \omega_{ci}^2 - \omega^2} - \frac{K_{\parallel}^2 \omega_{pi}^2}{K^2 \omega^2}$$

by using $\omega_{ci}^2 \gg \omega^2$

$$\chi_i = \frac{K_{\perp}^2 \omega_{pi}^2}{K^2 \omega_{ci}^2} - \frac{K_{\parallel}^2 \omega_{pi}^2}{K^2 \omega^2} \quad (21)$$

Similarly for χ_d , $F'_d = 1$ and $\omega_{cd} = 0$

So, the susceptibility for the un-magnetized and cold dust particles is obtained as

$$\chi_d = - \frac{\omega_{pd}^2}{\omega^2} \quad (22)$$

The dispersion relation for DLH wave in Quantum plasma will be

$$\varepsilon(\omega, \vec{K}) = 1 + \frac{1}{k^2 \lambda_{Fe}^2} + \frac{k_{\perp}^2 \omega_{pi}^2}{k^2 \omega_{ci}^2} - \frac{k_{\parallel}^2 \omega_{pi}^2}{k^2 \omega^2} - \frac{\omega_{pd}^2}{\omega^2} = 0$$

For

$$\begin{aligned} \frac{\omega_{pi}^2}{\omega_{ci}^2} &\gg \frac{1}{\omega^2 \lambda_{Fe}^2} \gg 1 \\ \omega^2 &= \frac{\omega_{dlh}^2 \left(1 + \frac{K_{\parallel}^2 \omega_{pi}^2}{K^2 \omega_{pd}^2} \right)}{\frac{k_{\perp}^2}{k^2} \left(\frac{\omega_{ci}^2}{c_{Fs}^2} \cdot \frac{1}{k_{\perp}^2} + 1 \right)} \end{aligned} \quad (23)$$

where

$$\omega_{dlh}^2 = \frac{\omega_{pd}^2 \omega_{ci}^2}{\omega_{pi}^2}, \quad C_{Fs}^{\prime 2} = \lambda_{Fe}^{\prime 2} \omega_{pi}^2 \quad \text{and} \quad \rho_{Fs}^{\prime 2} = \frac{C_{Fs}^{\prime 2}}{\omega_{ci}^2}$$

So

$$\omega^2 = \frac{\omega_{dlh}^2 \left(1 + \frac{K_{\parallel}^2 \omega_{pi}^2}{K^2 \omega_{pd}^2} \right)}{\frac{k_{\perp}^2}{k^2} \left(\frac{1}{k_{\perp}^2} \cdot \frac{1}{\rho_{Fs}^{\prime 2}} + 1 \right)} \quad (24)$$

Since

$$\begin{aligned} k_{\perp}^2 &\gg k_{\parallel}^2 \\ k_{\perp}^2 + k_{\parallel}^2 &= k^2 \\ k_{\perp}^2 &\sim k^2 \quad \Rightarrow \quad \frac{k_{\perp}^2}{k^2} = 1 \end{aligned}$$

Therefore

$$\omega^2 = \omega_{dlh}^2 \left(1 + \frac{K_{\parallel}^2 \omega_{pi}^2}{K^2 \omega_{pd}^2} \right) \times \left(1 + \frac{1}{K_{\perp}^2 \rho_{Fs}^{\prime 2}} \right)^{-1}$$

Or

$$\omega^2 = \omega_{dlh}^2 \left(1 + \frac{K_{\parallel}^2 \omega_{pi}^2}{K^2 \omega_{pd}^2} \right) \times \left(1 - \frac{1}{K_{\perp}^2 \rho_{Fs}^{\prime 2}} \right) \quad (25)$$

is the dispersion relation of the DLH wave in the cold Fermi dusty magneto plasma.

As

$$\lambda_{Fe}^{\prime 2} = \frac{V_{Fe}^2 (1 + \gamma_e)}{\omega_{pe}^2}$$

So, the relation for $C_{Fs}^{\prime 2}$ is

$$\begin{aligned} C_{Fs}^{\prime 2} &= \frac{\omega_{pi}^2}{\omega_{pe}^2} V_{pe}^2 (1 + \gamma_e) \\ C_{Fs}^{\prime 2} &= \frac{\omega_{pi}^2}{\omega_{pe}^2} \cdot \frac{2K_B T_{Fe}}{m_e} \left(1 + \frac{\hbar^2 K^2}{8 m_e K_B T_{Fe}} \right) \\ C_{Fs}^{\prime 2} &= \frac{2K_B T_{Fe}}{m_i} \left(1 + \frac{\hbar^2 K^2}{8 m_e K_B T_{Fe}} \right) \end{aligned}$$

3. Results and Discussion

- ρ_{Fs}^{\prime} is the ion gyro radius at the electron Fermi temperature with quantum correction as

$$\gamma_e = \frac{\hbar^2 k^2}{8 m_e k_B T_{Fe}}$$

- The DLH wave is seen to be significantly modified by the quantum effect.
 - In this case electrons at the Fermi temperature drive the wave.
- If the electrons and ions within the plasma have different temperatures such as $T_e \gg T_i$ then there is a sort of hybrid wave that depends upon the ion mass

m_i and Fermi temperature of electron T_{Fe} .




Figure 1. Variation of ω as a function of wave number k

Figure 1 shows that frequency of the quantum dust-lower-hybrid wave increases with the increase of k .




Figure 2. Variation of ω as a function of propagation angle θ

Figure 2 shows that frequency of the quantum dust-lower-hybrid wave increases at small angle of propagation.




Figure 3. Variation of ω as a function of external magnetic field B_0 and figure 3 shows that frequency of the quantum dust-lower-hybrid wave increases with the increase of magnetic field B_0 .

4. Conclusion

Equation (25) is the dispersion relation of the DLH wave in an ultra-cold and uniformly magnetized Fermi dusty plasma employing the quantum hydrodynamic model of a plasma with quantum and thermal corrections.

This relation is significantly affected by the quantum correction. This will find applications in diagnosing the charged dust impurities in microelectronics and wave particle interactions in the dusty quantum magnetoplasma. Bohm potential term shows how the concept of quantum potential leads to the notion of an “unbroken wholeness of entire universe”.

References

- [1] Haas F, Garcia LG, Goedert, J, Manfredi, G. Quantum ion-acoustic waves. *Phys. Plasmas*. 2003;10(10): 3858.
- [2] Markowich PA, Ringhofer CA, Schmeiser C. *Semiconductor Equations*. Vienna: Springer; 1990.
- [3] Kremp D, Bornath TH, Bonitz M, Schlanges M. Quantum kinetic theory of plasmas in strong laser fields. *Phys. Rev.* 1999;E 60; 4725.

- [4] Jung YD. Quantum-mechanical effects on electron-electron scattering in dense high-temperature plasmas. *Phys Plasmas*. 2001;8: 3842.
- [5] Opher M, Silva LO, Dager DE, Decyk VK, Dawson JM. Quantum ion-acoustic waves. *Phys Plasma*. 2001;8: 2454.
- [6] Chabrier G, Douchin F, Potekhin AY. Dense astrophysical plasmas. *J Phys: Condens Matter*. 2002;14: 9133.
- [7] Shukla PK. A new dust mode in quantum plasmas. *Phys Lett A*. 2006;352(3): 242–243.
- [8] Shukla PK, Stenflo L. Jeans instabilities in quantum dusty plasmas. *Phys Lett A*. 2006(4–5):355: 378–380.
- [9] Shukla PK, Ali S. Dust acoustic waves in quantum plasmas. *Phys Plasma*. 2005;12(11): 114502.
- [10] Shukla PK, Stenflo L, Bingham R. Shielding of a slowly moving test charge in a quantum plasmas. *Phys Lett A*. 2006;359(3): 218–219.